# UTILIZATION MANAGEMENT MEDICAL POLICY

**POLICY:** Oncology (Injectable) – Blincyto Utilization Management Medical Policy

• Blincyto® (blinatumomab intravenous infusion – Amgen)

**REVIEW DATE:** 09/10/2025

#### **OVERVIEW**

Blincyto, a bispecific CD19-directed CD3 T-cell engager, is indicated for the following uses:<sup>1</sup>

- Minimal residual disease (MRD)-positive, CD19-positive B-cell precursor acute lymphoblastic leukemia (ALL) in first or second complete remission with MRD ≥ 0.1% in patients ≥ 1 month of age.
- Relapsed or refractory CD19-positive B-cell ALL in patients  $\geq 1$  month of age.
- B-cell precursor, CD-19-positive Philadelphia chromosome-negative ALL in the consolidation phase of multiphase chemotherapy in patients ≥ 1 month of age.

## **Dosing Information**

Dosing in MRD-Positive B-Cell Precursor ALL. For patients > 45 kg (99 lbs), the dose of Blincyto is 28 mcg/day on Days 1 through 28 of each 42-day cycle. For patients < 45 kg (99 lbs), the dose is 15 mcg/m<sup>2</sup>/day, not to exceed 28 mcg/day on Days 1 through 28 of each 42-day cycle. A maximum of 4 cycles of Blincyto is recommended for MRD positive B-cell precursor ALL. A course of treatment consists of 1 induction cycle followed by 3 consolidation cycles. A treatment course may take 6 to 9 months to complete. Dosing in Relapsed/Refractory B-Cell Precursor ALL. For patients < 45 kg (99 lbs), Blincyto is dosed based on body surface area. The recommended dose in Cycle 1 is 5 mcg/m<sup>2</sup>/day (not to exceed 9 mcg/day) on Days 1 through 7 and 15 mcg/m<sup>2</sup>/day (not to exceed 28 mcg/day) on Days 8 through 28. In subsequent cycles, the recommended dose is 15 mcg/m<sup>2</sup>/day (not to exceed 28 mcg/day) on Days 1 through 28. For patients > 45 kg (99 lbs), the recommended dose in Cycle 1 is 9 mcg/day on Days 1 through 7 and 28 mcg/day on Days 8 through 28. In subsequent cycles, the recommended dose is 28 mcg/day on Days 1 through 28. A maximum of 9 cycles of Blincyto is recommended for relapsed/refractory B-cell precursor ALL. A treatment course of Blincyto consists of up to 2 induction cycles, 3 consolidation cycles, and up to 4 additional cycles. A cycle of induction or consolidation therapy consists of a 28-day continuous intravenous infusion followed by 14-day treatment-free interval. A single course of continued therapy consists of a 28-day continuous intravenous infusion followed by 56-day treatment-free interval.

<u>Dosing in B-Cell Precursor ALL in the Consolidation Phase</u>. For patients  $\geq$  45 kg (99 lbs), the dose of Blincyto is 28 mcg/day on Days 1 through 28 of each 42-day cycle. For patients < 45 kg (99 lbs), the dose is 15 mcg/m²/day, not to exceed 28 mcg/day on Days 1 through 28 of each 42-day cycle. Blincyto is given in cycles 1 and 2, then chemotherapy alone in cycles 3 – 5, then Blincyto in cycle 6, chemotherapy in cycle 7, and Blincyto in cycle 8.

#### Guidelines

The National Comprehensive Cancer Network (NCCN) guidelines on **Acute Lymphoblastic Leukemia** (version 2.2025 – June 27, 2025) and **Pediatric Acute Lymphoblastic Leukemia** (version 1.2026 – August 11, 2025) recommend Blincyto for relapsed/refractory B-cell ALL, induction therapy, consolidation therapy, and maintenance therapy.<sup>2-4</sup>

Safety

Oncology (Injectable) – Blincyto UM Medical Policy Page 2

Blincyto contains a boxed warning for cytokine release syndrome which may be life-threatening or fatal and neurologic toxicities which may be severe, life-threatening, or fatal. Stop or discontinue Blincyto as recommended for either toxicity.

#### **POLICY STATEMENT**

Prior Authorization is recommended for medical benefit coverage of Blincyto. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Extended approvals are allowed if the patient continues to meet the Criteria and Dosing. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Blincyto, as well as the monitoring required for adverse events and long-term efficacy, approval requires Blincyto to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

## RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Blincyto is recommended in those who meet the following criteria:

### **FDA-Approved Indication**

- 1. Acute Lymphoblastic Leukemia. Approve for 1 year if the patient meets BOTH of the following (A and B):
  - A) Patient has B-cell precursor disease; AND
  - **B)** The medication is prescribed by or in consultation with an oncologist.

**Dosing.** Approve up to 28 mcg/day administered by intravenous infusion on Days 1 through 28 of each treatment cycle with a minimum of a 14-day treatment-free interval between cycles.<sup>1</sup>

## CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Blincyto is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

#### REFERENCES

- 1. Blincyto® intravenous infusion [prescribing information]. Thousand Oaks, CA: Amgen; April 2025.
- 2. The NCCN Pediatric Acute Lymphoblastic Leukemia Oncology Guidelines (version 1.2026 August 11, 2025). © 2025 National Comprehensive Cancer Network. Available at: <a href="http://www.nccn.org">http://www.nccn.org</a>. Accessed on August 21, 2025.
- 3. The NCCN Acute Lymphoblastic Leukemia Clinical Practice Guidelines in Oncology (version 2.2025 June 27, 2025). © 2025 National Comprehensive Cancer Network. Available at: <a href="http://www.nccn.org">http://www.nccn.org</a>. Accessed on August 21, 2025.
- 4. The NCCN Drugs and Biologics Compendium. © 2025 National Comprehensive Cancer Network. Available at: <a href="http://www.nccn.org">http://www.nccn.org</a>. Accessed on August 21, 2025. Search term: blinatumomab.

# HISTORY

Type of Revision	Summary of Changes	Review Date

# Oncology (Injectable) – Blincyto UM Medical Policy Page 3

Annual Revision	Acute Lymphoblastic Leukemia: For Philadelphia chromosome positive disease,	09/06/2023
	patient has relapsed or refractory disease and medication is used for maintenance	
	therapy were added as additional options for approval.	
Annual Revision	Acute Lymphoblastic Leukemia: For Philadelphia chromosome negative disease,	09/04/2024
	removed patient is minimal residual disease positive as an option for approval. Added	
	the medication is used for induction therapy as an option for approval. For Philadelphia	
	chromosome-like disease removed minimal residual disease positive and added the	
	medication is used for consolidation therapy. For Philadelphia chromosome-positive	
	disease added the medication is used for induction therapy as an option for approval.	
	Removed the patient has tried at least one tyrosine kinase inhibitor, the patient does	
	not have a complete response to induction therapy, and patient is minimal residual	
	disease positive as options for approval.	
Annual Revision	Acute Lymphoblastic Leukemia: The requirements that the patient is Philadelphia	09/10/2025
	chromosome negative, Philadelphia chromosome-like, or Philadelphia chromosome	
	positive were removed. The specifications around treatment phases such as induction,	
	consolidation, maintenance, and relapsed/refractory disease were removed.	