Opdivo Qvantig™ (nivolumab and hyaluronidase-nvhy) (Subcutaneous)

Document Number: IC-0784

Last Review Date: 08/04/2026
Date of Origin: 02/04/2025
Dates Reviewed: 02/2025, 10/2025, 12/2025, 02/2026, 05/2026, 08/2026

I. Length of Authorization Δ1

- Initial: Prior authorization validity will be provided initially for 6 months (180 days), unless otherwise specified.
  - Substitution/Switch-Therapy for Intravenous Nivolumab: Prior authorization validity will follow the same parameters for intravenous nivolumab based on indication requested *[see Opdivo IV policy – Document Number: IC-0226]*
- Renewal: Prior authorization validity may be renewed every 6 months (180 days) thereafter, unless otherwise specified.
  - Substitution/Switch-Therapy for Intravenous Nivolumab: Prior authorization validity will follow the same parameters for intravenous nivolumab based on indication requested *[see Opdivo IV policy – Document Number: IC-0226]*
  - NSCLC neoadjuvant treatment: Prior authorization validity may NOT be renewed.
  - NSCLC neoadjuvant treatment followed by adjuvant treatment: Prior authorization validity may be provided for a maximum of 4 neoadjuvant doses and 13 adjuvant doses.
  - Prior authorization validity may be renewed up to a maximum of 1 year of therapy* for the following:
    - Cutaneous Melanoma (adjuvant therapy)
    - Esophageal Cancer (adjuvant treatment of resected esophageal or gastroesophageal junction cancer)
    - Urothelial Carcinoma (adjuvant therapy)
  - Prior authorization validity may be renewed up to a maximum of 2 years of therapy* for the following:
    - Esophageal Cancer (first-line therapy squamous cell carcinoma)
    - Gastric Cancer, Gastroesophageal Junction Cancer, and Esophageal Adenocarcinoma
    - Hepatocellular Carcinoma (first-line therapy)
    - Renal Cell Carcinoma (in combination with cabozantinib)
    - Urothelial Carcinoma (first line therapy in combination with gemcitabine and cisplatin, followed by single-agent maintenance therapy)

© 2026 Prime Therapeutics Management, LLC
Proprietary & Confidential.

- ❖ Colorectal Cancer (following combination treatment with intravenous nivolumab/ipilimumab)

Note: The maximum number of doses is dependent on the dosing frequency and duration of therapy. Refer to Section V for exact dosage.		
Dosing Frequency	**Maximum length of therapy**	**Maximum number of doses**
2 weeks	1 year	26 doses
	2 years	52 doses
3 weeks	2 years	35 doses
4 weeks	1 year	13 doses
	2 years	26 doses

II. Dosing Limits

Max Units (per dose and over time) [HCPCS Unit]:

- Renal Cell Carcinoma, HCC, SCCHN, Cutaneous Melanoma, CRC: 600 billable units every 28 days
- All Other Indications: 1800 billable units every 84 days

III. Initial Approval Criteria [1]

Prior authorization validity is provided in the following conditions:

- Member is at least 18 years of age, unless otherwise specified; **AND**

Universal Criteria

- Member has not received previous therapy with a programmed death (PD-1/PD-L1)-directed therapy, unless otherwise specified [Δ] *(Note: Not applicable when used as switch-therapy from intravenous nivolumab)*; **AND**
- Therapy will not be used concomitantly with intravenous nivolumab; **AND**
- Therapy will not be used concurrently with intravenous chemotherapy agents *(not applicable when used for FDA approved combination therapy indicated with* **†***)*; **AND**
- Intravenous nivolumab must be used in the following:
  - Members < 80 kg; **OR**
  - Members receiving therapy in combination with ipilimumab; **AND**

Substitution/Switch-Therapy for Intravenous Nivolumab [Δ] ‡

- Used as substitution for OR switch-therapy from intravenous nivolumab; **AND**
  - Member has previously met criteria for use of intravenous nivolumab *[see Opdivo IV policy – Document Number: IC-0226];* **AND**
    - Member has been receiving treatment with intravenous nivolumab and has shown a beneficial disease response and absence of unacceptable toxicity while on therapy; **OR**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

- Member currently meets criteria for use of intravenous nivolumab *[see Opdivo IV policy – Document Number: IC-0226]*

Urothelial Carcinoma † [1]

- Used as a single agent; **AND**
  - Used for locally advanced or metastatic disease that progressed during or following platinum-containing chemotherapy OR have disease progression within 12 months of neoadjuvant or adjuvant treatment with a platinum-containing regimen; **OR**
  - Used as adjuvant therapy in members who are at a high risk for disease recurrence** after undergoing radical surgical resection; **OR**
- Used in combination with cisplatin and gemcitabine, followed by single agent maintenance therapy; **AND**
  - Used as first line therapy in members with unresectable or metastatic disease

***** Note:*** [1,62]

– *High risk for disease recurrence is defined as:*

- *ypT2-ypT4a or ypN+ for members who received neoadjuvant platinum-based therapy (excluding urothelial carcinoma of the prostate with stromal invasion); **OR***
- *pT3-pT4a or pN+ for members who did not receive neoadjuvant platinum-based therapy*

Colorectal Cancer (CRC) † [1]

- Member has microsatellite instability-high (MSI-H)/mismatch repair deficient (dMMR) disease as determined by an FDA-approved or Clinical Laboratory Improvement Amendments (CLIA) compliant test❖; **AND**
- Used as a single agent; **AND**
- Member is at least 12 years of age; **AND**
  - Used for unresectable or metastatic disease following combination treatment with intravenous nivolumab/ipilimumab; **OR**
  - Used as subsequent therapy for metastatic disease that has progressed following treatment with a fluoropyrimidine, oxaliplatin and irinotecan regimen

Esophageal Cancer † [1]

- Used as a single agent; **AND**
  - Used as adjuvant treatment of completely resected esophageal or gastroesophageal junction (GEJ) cancer with residual pathologic disease in members who have received neoadjuvant chemoradiotherapy (CRT); **OR**
  - Used as subsequent therapy after prior fluoropyrimidine- and platinum-based chemotherapy; **AND**
    - Used for unresectable advanced, recurrent, or metastatic esophageal squamous cell carcinoma (ESCC); **OR**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

- Used in combination with fluoropyrimidine- and platinum-containing chemotherapy; **AND**
  - Used as first-line therapy; **AND**
  - Tumor expresses PD-L1 (CPS ≥1) as determined by an FDA-approved or CLIA compliant test❖; **AND**
  - Used in members with unresectable, advanced or metastatic ESCC

Gastric Cancer, Gastroesophageal Junction Cancer, and Esophageal Adenocarcinoma † [1]

- Used in combination with fluoropyrimidine- and platinum-containing chemotherapy; **AND**
- Used as first-line therapy; **AND**
- Member has advanced or metastatic gastric cancer, gastroesophageal junction cancer, or esophageal adenocarcinoma; **AND**
- Tumor expresses PD-L1 (CPS ≥1) as determined by an FDA-approved or CLIA compliant test❖

Squamous Cell Carcinoma of the Head and Neck (SCCHN) † [1]

- Used as single-agent therapy; **AND**
- Member has recurrent or metastatic disease with disease progression on or after platinum-based therapy; **AND**
- Member does not have disease of the nasopharynx

Hepatocellular Carcinoma (HCC) † [1]

- Used as single agent therapy; **AND**
- Member has unresectable or metastatic disease; **AND**
  - Used as first-line therapy following combination treatment with intravenous nivolumab/ipilimumab; **OR**
  - Used as subsequent therapy after being previously treated with sorafenib and following combination treatment with intravenous nivolumab/ipilimumab

Renal Cell Carcinoma (RCC) † [1]

- Used as a single agent; **AND**
  - Used as first line therapy in members with intermediate or poor risk advanced disease following treatment with intravenous nivolumab/ipilimumab; **OR**
  - Used as subsequent therapy for advanced disease after prior anti-angiogenic therapy; **OR**
- Used in combination with cabozantinib (Cabometyx only); **AND**
  - Used as first-line therapy for advanced disease

Cutaneous Melanoma † [1]

- Used as single agent therapy; **AND**
- Member is at least 12 years of age; **AND**

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

- Used as first-line therapy for unresectable or metastatic disease; **OR**
- Used as subsequent therapy for unresectable or metastatic disease following treatment with intravenous nivolumab/ipilimumab; **OR**
- Used as adjuvant treatment and member has stage IIB, stage IIC, stage III or metastatic disease and has undergone complete resection

Non-Small Cell Lung Cancer (NSCLC) † [1]

- Used as single-agent therapy; **AND**
  - Used for metastatic disease; **AND**
  - Used as subsequent therapy after progression on or after platinum-based chemotherapy (Note: Members with EGFR or ALK genomic tumor aberrations should have disease progression on targeted therapies prior to receiving subcutaneous nivolumab); **OR**
- Used in combination with platinum-doublet chemotherapy for resectable (tumors ≥ 4 cm or node positive) disease; **AND**
  - Used as neoadjuvant therapy; **OR**
  - Used as neoadjuvant therapy followed by single-agent subcutaneous nivolumab therapy as adjuvant treatment after surgery; **AND**
    - Member has no known EGFR mutations or ALK rearrangements

❖*If confirmed using an FDA approved assay – http://www.fda.gov/companiondiagnostics*

† FDA Approved Indication(s); **‡** Compendia Recommended Indication(s); **Φ** Orphan Drug

IV. Renewal Criteria [Δ 1]

Prior authorization validity may be renewed based upon the following criteria:

- Member continues to meet universal and other indication-specific relevant criteria such as concomitant therapy requirements (not including prerequisite therapy), performance status, etc. identified in section III; **AND**
- Duration of authorization has not been exceeded *(refer to Section I)*; **AND**
- Disease response with treatment as defined by stabilization of disease or decrease in size of tumor or tumor spread; **AND**
- Absence of unacceptable toxicity from the drug. Examples of unacceptable toxicity include: severe immune-mediated adverse reactions (e.g., pneumonitis, hepatitis, colitis, endocrinopathies, nephritis/renal dysfunction, rash/dermatitis [including Stevens-Johnson syndrome (SJS), drug rash with eosinophilia and systemic symptoms (DRESS), and toxic epidermal necrolysis (TEN)], myocarditis, pericarditis, vasculitis, solid organ transplant rejection, etc.), complications of allogeneic hematopoietic stem cell transplantation (HSCT), etc.

Δ Notes:

– Members responding to therapy who relapse ≥ 6 months after discontinuation due to duration are

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

eligible to re-initiate PD-directed therapy.

- Members previously presenting with aggressive disease who are exhibiting stable disease on treatment as their best response (or if therapy improved performance status) may be eligible for continued therapy without interruption or discontinuation.
- Members who complete adjuvant therapy and progress ≥ 6 months after discontinuation are eligible to re-initiate PD-directed therapy for metastatic disease.
- Members whose tumors, upon re-biopsy, demonstrate a change in actionable mutation (e.g., MSS initial biopsy; MSI-H subsequent biopsy) may be eligible to re-initiate PD-directed therapy and will be evaluated on a case-by-case basis.

V. Dosage/Administration Δ 1

Indication	Dose
Substitution/Switch-Therapy for Intravenous Nivolumab ‡	– Dose does not exceed one of the following: ○ 600 mg/10,000 units every 2 weeks ○ 900 mg/15,000 units every 3 weeks ○ 1,200 mg/20,000 units every 4 weeks; **AND** – Dosing will follow the same indication-specific parameters for IV nivolumab *[see Opdivo IV policy – Document Number: IC-0226]*
Renal Cell Carcinoma †	**Single agent:** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity; **OR** **Combination with cabozantinib:** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity for up to 2 years
Cutaneous Melanoma †	**Unresectable or metastatic disease:** – Adult and pediatric members 12 years of age and older and weighing 40 kg or greater♦: 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity – Pediatric members 12 years of age and older and weighing 30 kg or greater but less than 40 kg♦: 300 mg/5,000 units every 2 weeks or 600 mg/10,000 units every 4 weeks, until disease progression or unacceptable toxicity **Adjuvant treatment:** – Adult and pediatric members 12 years of age and older and weighing 40 kg or greater♦: 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks until disease recurrence or unacceptable toxicity for up to 1 year – Pediatric members 12 years of age and older and weighing 30 kg or greater but less than 40 kg♦: 300 mg/5,000 units every 2 weeks or 600 mg/10,000 units every 4 weeks until disease recurrence or unacceptable toxicity for up to 1 year
Non-Small Cell Lung Cancer (NSCLC) †	**Neoadjuvant treatment:** – 900 mg/15,000 units with platinum-doublet chemotherapy on the same day every 3 weeks for up to 3 cycles

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

	Neoadjuvant treatment followed by adjuvant treatment: – 900 mg/15,000 units with platinum-doublet chemotherapy on the same day every 3 weeks for up to 4 cycles, followed by single-agent adjuvant treatment after surgery at 1,200 mg/20,000 units every 4 weeks until disease progression, recurrence, or unacceptable toxicity for up to 13 cycles (up to one year) **Metastatic disease:** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity
Urothelial Carcinoma †	**Adjuvant treatment:** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease recurrence or unacceptable toxicity for up to 1 year **Disease progression:** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity **First-line therapy (in combination with gemcitabine and cisplatin, followed by single agent maintenance therapy):** – 900 mg/15,000 units every 3 weeks with cisplatin and gemcitabine on the same day for up to 6 cycles, followed by: – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity for up to 2 years
Esophageal Cancer †	**First line therapy (squamous cell carcinoma):** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, in combination with fluoropyrimidine- and platinum-containing chemotherapy every 2 weeks, until disease progression or unacceptable toxicity for up to 2 years **Subsequent therapy (squamous cell carcinoma):** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity **Adjuvant treatment of resected esophageal or gastroesophageal junction cancer:** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease recurrence or unacceptable toxicity for up to 1 year
Gastric Cancer, Gastroesophageal Junction Cancer, and Esophageal Adenocarcinoma †	– 600 mg/10,000 units every 2 weeks in combination with fluoropyrimidine- and platinum-containing chemotherapy every 2 weeks, until disease progression or unacceptable toxicity for up to 2 years; **OR** – 900 mg/15,000 units every 3 weeks with fluoropyrimidine- and platinum containing chemotherapy every 3 weeks, until disease progression or unacceptable toxicity for up to 2 years
Hepatocellular Carcinoma (HCC) †	**First line therapy:** – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity for up to 2 years

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

	Subsequent therapy: – 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity
Colorectal Cancer (CRC) †	**Following progression on fluoropyrimidine, oxaliplatin and irinotecan regimen:** – Adult and pediatric members 12 years of age and older and weighing 40 kg or greater♦: 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity – Pediatric members 12 years of age and older and weighing 30 kg or greater but less than 40 kg♦: 300 mg/5,000 units every 2 weeks or 600 mg/10,000 units every 4 weeks, until disease progression or unacceptable toxicity **Following combination treatment with intravenous nivolumab/ipilimumab:** – Adult and pediatric members 12 years of age and older and weighing 40 kg or greater♦: 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity for up to 2 years – Pediatric members 12 years of age and older and weighing 30 kg or greater but less than 40 kg♦: 300 mg/5,000 units every 2 weeks or 600 mg/10,000 units every 4 weeks, until disease progression or unacceptable toxicity for up to 2 years
Squamous Cell Carcinoma of the Head and Neck (SCCHN) †	– 600 mg/10,000 units every 2 weeks or 1,200 mg/20,000 units every 4 weeks, until disease progression or unacceptable toxicity
Note: – *♦Dosing for adult and pediatric members weighing 40 kg or greater AND for pediatric members weighing 30 kg or greater but less than 40 kg is listed in the prescribing information; however, the IV formulation of nivolumab should be used for members weighing < 80 kg.* – *Opdivo Qvantig (nivolumab and hyaluronidase-nvhy) has different dosage and administration instructions than intravenous nivolumab products.* – *Opdivo Qvantig is for subcutaneous use only in the abdomen or thigh.* – *Opdivo Qvantig is to be administered by a healthcare professional only.* – *Opdivo Qvantig is for subcutaneous use only administered over 3-5 minutes.*	

VI. Billing Code/Availability Information

HCPCS Code:

- J9289 – Injection, nivolumab, 2 mg and hyaluronidase-nvhy; 1 billable unit = 2 mg

NDC(s):

- Opdivo Qvantig single-dose vial providing 300 mg nivolumab and 5,000 units hyaluronidase per 2.5 mL (120 mg/2,000 units per mL): 00003-3120-xx
- Opdivo Qvantig single-dose vial providing 600 mg nivolumab and 10,000 units hyaluronidase per 5 mL (120 mg/2,000 units per mL): 00003-6120-xx

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

VII. References

1. Opdivo Qvantig [package insert]. Princeton, NJ; Bristol-Myers Squibb, Inc; May 2026. Accessed July 2026.
2. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) nivolumab and hyaluronidase-nvhy. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
3. Scherpereel A, Mazieres J, Greillier L, et al. Second- or third-line nivolumab (Nivo) versus nivo plus ipilimumab (Ipi) in malignant pleural mesothelioma (MPM) patients: Results of the IFCT-1501 MAPS2 randomized phase II trial. [Abstract]. J Clin Oncol 2017;35: Abstract LBA 8507.
4. Walocko FM, Scheier BY, Harms PW, et al. Metastatic Merkel cell carcinoma response to nivolumab. J Immunother Cancer. 2016 Nov 15;4:79.
5. Tawbi HA, Forsyth PA, Algazi A, et al. Combined Nivolumab and Ipilimumab in Melanoma Metastatic to the Brain. N Engl J Med. 2018 Aug 23;379(8):722-730.
6. Morris VK, Salem ME, Nimeiri H, et al. Nivolumab for previously treated unresectable metastatic anal cancer (NCI9673): a multicentre, single-arm, phase 2 study. Lancet Oncol. 2017 Apr;18(4):446-453. Doi: 10.1016/S1470-2045(17)30104-3. Epub 2017 Feb 18.
7. Zhao X, Ivaturi V, Gopalakrishnan M, et al. Abstract CT 101: A model-based exposure-response (E-R) assessment of a nivolumab (NIVO) 4-weekly (Q4W) dosing schedule across multiple tumor types. Cancer Res July 1 2017 (77) (13 Supplement) CT101; DOI**:** 10.1158/1538-7445.AM2017-CT101.
8. Zhao X, Suryawanshi M, Hruska M, et al. Assessment of nivolumab benefit-risk profile of a 240 mg flat dose relative to a 3 mg/kg dosing regimen in patients with advanced tumors. Ann Oncol2017; 28:2002-2008.
9. Feng Y, Xiaoning W, Bajaj G, et al. Nivolumab exposure-response analyses of efficacy and safety in previously treated squamous or nonsquamous non-small cell lung cancer. ClinCa Res 2017;23(18): 5394-5405.
10. Gupta S, Bellmunt J, Plimack ER, et al. Defining “platinum-ineligible” patients with metastatic urothelial cancer (mUC). J Clin Oncol. 2022 June 1;40(16_suppl):4577.
11. Hellmann MD, Ciuleanu TE, Pluzanski A, et al. Nivolumab plus ipilimumab in lung cancer with a high tumor mutational burden. N Engl J Med 2018; 378:2093-2104.
12. Fahrenbruch R, Kintzel P, Bott AM, et al. Dose Rounding of Biologic and Cytotoxic Anticancer Agents: A Position Statement of the Hematology/Oncology Pharmacy Association. J Oncol Pract. 2018 Mar;14(3):e130-e136.
13. Hematology/Oncology Pharmacy Association. *Intravenous Cancer Drug Waste Issue Brief*. Updated February 2024. Available at https://www.hoparx.org/documents/287/HOPA_Drug_Waste_Issue_Brief_-_Updated_02.22.24.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

14. Bach PB, Conti RM, Muller RJ, et al. Overspending driven by oversized single dose vials of cancer drugs. BMJ. 2016 Feb 29;352:i788.
15. Weber JS, D'Angelo SP, Minor D, et al. Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. Lancet Oncol. 2015 Apr;16(4):375-84. Doi: 10.1016/S1470-2045(15)70076-8. Epub 2015 Mar 18.
16. Robert C, Long GV, Brady B, et al. Nivolumab in previously untreated melanoma without BRAF mutation. N Engl J Med. 2015 Jan 22;372(4):320-30. Doi: 10.1056/NEJMoa1412082. Epub 2014 Nov 16.
17. Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. N Engl J Med. 2015 Jul 2;373(1):23-34. Doi: 10.1056/NEJMoa1504030. Epub 2015 May 31.
18. Weber J, Mandala M, Del Vecchio M, et al. Adjuvant Nivolumab versus Ipilimumab in Resected Stage III or IV Melanoma. N Engl J Med. 2017 Nov 9;377(19):1824-1835. Doi: 10.1056/NEJMoa1709030. Epub 2017 Sep 10.
19. Algazi AP, Tsai KK, Shoushtari AN, et al. Clinical outcomes in metastatic uveal melanoma treated with PD-1 and PD-L1 antibodies. Cancer. 2016 Nov 15;122(21):3344-3353. Doi: 10.1002/cncr.30258. Epub 2016 Aug 17.
20. Piulats JM, Espinosa E, de la Cruz-Merino L, et al. Nivolumab plus ipilimumab for treatment-naive metastatic uveal melanoma: An open-label, multicenter, phase II trial by the Spanish Multidisciplinary Melanoma Group (GEM-1402). J Clin Oncol 2021; 39:586-598.
21. El-Khoueiry AB, Sangro B, Yau T, et al. Nivolumab in patients with advanced hepatocellular carcinoma (CheckMate 040): an open-label, non-comparative, phase ½ dose escalation and expansion trial. Lancet. 2017 Jun 24;389(10088):2492-2502. Doi: 10.1016/S0140-6736(17)31046-2. Epub 2017 Apr 20.
22. Brahmer J, Reckamp KL, Baas P, et al. Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer. N Engl J Med. 2015 Jul 9;373(2):123-35. Doi: 10.1056/NEJMoa1504627. Epub 2015 May 31.
23. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer. N Engl J Med. 2015 Oct 22;373(17):1627-39. Doi: 10.1056/NEJMoa1507643. Epub 2015 Sep 27.
24. Antonia SJ, López-Martin JA, Bendell J, et al. Nivolumab alone and nivolumab plus ipilimumab in recurrent small-cell lung cancer (CheckMate 032): a multicentre, open-label, phase ½ trial. Lancet Oncol. 2016 Jul;17(7):883-895. Doi: 10.1016/S1470-2045(16)30098-5. Epub 2016 Jun 4.
25. Motzer RJ, Escudier B, McDermott DF, et al. Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma. N Engl J Med. 2015 Nov 5;373(19):1803-13. Doi: 10.1056/NEJMoa1510665. Epub 2015 Sep 25.
26. Motzer RJ, Tannir NM, McDermott DF, et al. Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma. N Engl J Med. 2018 Apr 5;378(14):1277-1290. Doi: 10.1056/NEJMoa1712126. Epub 2018 Mar 21.
27. Armand P, Engert A, Younes A, et al. Nivolumab for Relapsed/Refractory Classic Hodgkin Lymphoma After Failure of Autologous Hematopoietic Cell Transplantation: Extended Follow-Up

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

of the Multicohort Single-Arm Phase II CheckMate 205 Trial. J Clin Oncol. 2018 May 10;36(14):1428-1439. Doi: 10.1200/JCO.2017.76.0793. Epub 2018 Mar 27.
28. Ansell SM, Lesokhin AM, Borrello I, et al. PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma. N Engl J Med. 2015 Jan 22;372(4):311-9. Doi: 10.1056/NEJMoa1411087. Epub 2014 Dec 6.
29. Ferris RL, Blumenschein G Jr, Fayette J, et al. Nivolumab for Recurrent Squamous-Cell Carcinoma of the Head and Neck. N Engl J Med. 2016 Nov 10;375(19):1856-1867. Epub 2016 Oct 8.
30. Sharma P, Retz M, Siefker-Radtke A, et al. Nivolumab in metastatic urothelial carcinoma after platinum therapy (CheckMate 275): a multicentre, single-arm, phase 2 trial. Lancet Oncol. 2017 Mar;18(3):312-322. Doi: 10.1016/S1470-2045(17)30065-7. Epub 2017 Jan 26.
31. Overman MJ, McDermott R, Leach JL, et al. Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study. Lancet Oncol. 2017 Sep;18(9):1182-1191. Doi: 10.1016/S1470-2045(17)30422-9. Epub 2017 Jul 19.
32. Overman MJ, Lonardi S, Wong KYM, et al. Durable Clinical Benefit With Nivolumab Plus Ipilimumab in DNA Mismatch Repair-Deficient/Microsatellite Instability-High Metastatic Colorectal Cancer. J Clin Oncol. 2018 Mar 10;36(8):773-779. Doi: 10.1200/JCO.2017.76.9901. Epub 2018 Jan 20.
33. Topalian SL, Bhatia S, Hollebecque A, et al. Non-comparative, open-label, multiple cohort, phase ½ study to evaluate nivolumab (NIVO) in patients with virus-associated tumors (CheckMate 358): Efficacy and safety in Merkel cell carcinoma (MCC). DOI: 10.1158/1538-7445.AM2017-CT074 Published July 2017.
34. Long GV, Atkinson V, Lo S, et al. Combination nivolumab and ipilimumab or nivolumab alone in melanoma brain metastases: a multicentre randomised phase 2 study. Lancet Oncol. 2018 May;19(5):672-681. Doi: 10.1016/S1470-2045(18)30139-6. Epub 2018 Mar 27.
35. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Anal Carcinoma. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
36. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Gestational Trophoblastic Neoplasia. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
37. Scherpereel A, Mazieres J, Greillier L, et al. Nivolumab or nivolumab plus ipilimumab in patients with relapsed malignant pleural mesothelioma (IFCT-1501 MAPS2): a multicentre, open-label, randomised, non-comparative, phase 2 trial. Lancet Oncol. 2019 Feb;20(2):239-253. Doi: 10.1016/S1470-2045(18)30765-4. Epub 2019 Jan 16.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

38. Disselhorst MJ, Quispel-Janssen J, Lalezari F, et al. Ipilimumab and nivolumab in the treatment of recurrent malignant pleural mesothelioma (INITIATE): results of a prospective, single-arm, phase 2 trial. Lancet Respir Med. 2019 Mar;7(3):260-270. Doi: 10.1016/S2213-2600(18)30420-X. Epub 2019 Jan 16.
39. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Small Bowel Adenocarcinoma. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
40. Chan TSY, Li J, Loong F, et al. PD1 blockade with low-dose nivolumab in NK/T cell lymphoma failing L-asparaginase: efficacy and safety. Ann Hematol. 2018 Jan;97(1):193-196. Doi: 10.1007/s00277-017-3127-2. Epub 2017 Sep 6.
41. Goldman JW, Crino L, Vokes EE, et al. P2.36: Nivolumab (nivo) in patients (pts) with advanced (adv) NSCLC and central nervous system (CNS) metastases (mets). J Thorac Oncol 2016;11:S238-S239.
42. Gauvain C, Vauleon E, Chouaid C, et al. Intracerebral efficacy and tolerance of nivolumab in non–small-cell lung cancer patients with brain metastases. Lung Cancer. 2018 Feb; 116:62-66. Doi: 10.1016/j.lungcan.2017.12.008.
43. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Non-Small Cell Lung Cancer. Version 6.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
44. Kato K, Cho BC, Takahashi M, et al. Nivolumab versus chemotherapy in patients with advanced esophageal squamous cell carcinoma refractory or intolerant to previous chemotherapy (ATTRACTION-3): a multicentre, randomised, open-label, phase 3 trial. Lancet Oncol. 2019;20(11):1506-1517. Doi:10.1016/S1470-2045(19)30626-6.
45. Hellmann MD, Paz-Ares L, Bernabe Caro R, et al. Nivolumab plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer. N Engl J Med. 2019;381(21):2020-2031. Doi:10.1056/NEJMoa1910231.
46. Reck M, Ciuleanu T-E, Dols MC, et al. Nivolumab (NIVO) + ipilimumab (IPI) + 2 cycles of platinum-doublet chemotherapy (chemo) vs 4 cycles chemo as first-line (1L) treatment (tx) for stage IV/recurrent non-small cell lung cancer (NSCLC): CheckMate 9LA [abstract]. J Clin Oncol 2020;38:Abstract 9501-9501.
47. Zalcman G, Peters S, Mansfield AS, et al. Checkmate 743: A phase 3, randomized, open-label trial of nivolumab (nivo) plus ipilimumab (ipi) vs pemetrexed plus cisplatin or carboplatin as first-line therapy in unresectable pleural mesothelioma. Journal of Clinical Oncology 2017 35:15_suppl, TPS8581-TPS8581

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

48. Azad NS, Gray RJ, Overman MJ, et al. Nivolumab Is Effective in Mismatch Repair-Deficient Noncolorectal Cancers: Results From Arm Z1D-A Subprotocol of the NCI-MATCH (EAY131) Study. J Clin Oncol. 2020 Jan 20;38(3):214-222.
49. Naumann RW, Hollebecque A, Meyer T, et al. Safety and Efficacy of Nivolumab Monotherapy in Recurrent or Metastatic Cervical, Vaginal, or Vulvar Carcinoma: Results From the Phase I/II CheckMate 358 Trial. J Clin Oncol. 2019 Nov 1;37(31):2825-2834.
50. Choueiri TK, Powles T, Burotto M, et al. 696O_PR Nivolumab + cabozantinib vs sunitinib in first-line treatment for advanced renal cell carcinoma: First results from the randomized phase III CheckMate 9ER trial. Volume 31, SUPPLEMENT 4, S1159, September 01, 2020.
51. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Bladder Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
52. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Esophageal and Esophagogastric Junction Cancers. Version 3.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
53. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Gastric Cancer. Version 3.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
54. Herrera AF, Moskowitz AJ, Bartlett NL, et al. Interim results of brentuximab vedotin in combination with nivolumab in patients with relapsed or refractory Hodgkin lymphoma. Blood. 2018 Mar 15;131 (11):1183-1194.
55. Cole PD, Mauz-Körholz C, Mascarin M, et al. HL-032: Nivolumab and Brentuximab Vedotin (BV)–Based, Response-Adapted Treatment in Children, Adolescents, and Young Adults (CAYA) With Standard-Risk Relapsed/Refractory Classical Hodgkin Lymphoma (R/R cHL): Primary Analysis of the Standard-Risk Cohort of the Phase 2 CheckMate 744 Study. Clinical Lymphoma Myeloma and Leukemia. Volume 20, Supplement 1, September 2020, Pages S245-S246.
56. Moehler M, Shitara K, Garrido M, et al. Nivolumab (nivo) plus chemotherapy (chemo) versus chemo as first-line (1L) treatment for advanced gastric cancer/gastroesophageal junction cancer (GC/GEJC)/esophageal adenocarcinoma (EAC): First results of the CheckMate 649 study. [abstract]. Presented at the Oral Presentation presented at the ESMO 2020 Annual Meeting; September 19-21, 2020; Virtual Meeting.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

57. Kelly RJ, Ajani JA, Kuzdzal J, et al. Adjuvant Nivolumab in Resected Esophageal or Gastroesophageal Junction Cancer. N Engl J Med. 2021 Apr 1;384(13):1191-1203. Doi: 10.1056/NEJMoa2032125.
58. Nivolumab. Micromedex Solutions. Greenwood Village, CO: Truven Health Analytics. http://micromedex.com/. Updated September 09, 2025. Accessed September 2025.
59. Lenz HJ, Lonardi S, Zagonel V, et al. Nivolumab (NIVO) + low-dose ipilimumab (IPI) as first-line (1L) therapy in microsatellite instability-high/DNA mismatch repair deficient (MSI-H/dMMR) metastatic colorectal cancer (mCRC): Clinical update [abstract]. Journal of Clinical Oncology 2019;37:3521-3521.
60. Bellmunt, J, Valderrama BP (2026). Treatment of metastatic urothelial cancer of the bladder and urinary tract. In Lerner SP, Shah S (Eds.), *UptoDate.* Last updated Feb 05, 2026. Accessed July 8, 2026. Available from https://www.uptodate.com/contents/treatment-of-metastatic-urothelial-cancer-of-the-bladder-and-urinary-tract.
61. Ready NE, Ott PA, Hellmann MD, et al. Nivolumab Monotherapy and Nivolumab Plus Ipilimumab in Recurrent Small Cell Lung Cancer: Results From the CheckMate 032 Randomized Cohort. J Thorac Oncol. 2020 Mar;15(3):426-435. Doi: 10.1016/j.jtho.2019.10.004.
62. Bajorin DF, Witjes JA, Gschwend JE, et al. Adjuvant Nivolumab versus Placebo in Muscle-Invasive Urothelial Carcinoma. N Engl J Med. 2021 Jun 3;384(22):2102-2114. Doi: 10.1056/NEJMoa2034442.
63. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Cervical Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
64. Fennell DA, Ewings S, Ottensmeier C, et al. Nivolumab versus placebo in patients with relapsed malignant mesothelioma (CONFIRM): a multicentre, double-blind, randomised, phase 3 trial. Lancet Oncol 2021; 22:1530.
65. Topalian SL, Bhatia S, Amin A, et al. Neoadjuvant Nivolumab for Patients With Resectable Merkel Cell Carcinoma in the CheckMate 358 Trial. J Clin Oncol. 2020;38(22):2476-2487. Doi:10.1200/JCO.20.00201.
66. Forde PM, Spicer J, Lu S et al. Neoadjuvant Nivolumab plus Chemotherapy in Resectable Lung Cancer. N Engl J Med. 2022 May 26;386(21):1973-1985. doi: 10.1056/NEJMoa2202170. Epub 2022 Apr 11. PMID: 35403841; PMCID: PMC9844511.
67. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Mesothelioma: Peritoneal. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
68. Scherpereel A, Mazieres J, Greillier L, et al; French Cooperative Thoracic Intergroup. Nivolumab or nivolumab plus ipilimumab in patients with relapsed malignant pleural mesothelioma (IFCT-1501 MAPS2): a multicentre, open-label, randomised, non-comparative, phase 2 trial.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

Lancet Oncol. 2019 Feb;20(2):239-253. Doi: 10.1016/S1470-2045(18)30765-4. Epub 2019 Jan 16. Erratum in: Lancet Oncol. 2019 Mar;20(3):e132.
69. Doki Y, Ajani JA, Kato K, et al. Nivolumab Combination Therapy in Advanced Esophageal Squamous-Cell Carcinoma. N Engl J Med. 2022 Feb 3;386(5):449-462. Doi: 10.1056/NEJMoa2111380.
70. De Santis M, Bellmunt J, Mead G, et al. Randomized phase II/III trial assessing gemcitabine/ carboplatin and methotrexate/carboplatin/vinblastine in patients with advanced urothelial cancer "unfit" for cisplatin-based chemotherapy: phase II—results of EORTC study 30986. J Clin Oncol. 2009 Nov 20;27(33):5634-9. Doi: 10.1200/JCO.2008.21.4924. Epub 2009 Sep 28.
71. Bouffet E, Larouche V, Campbell BB, et al. Immune Checkpoint Inhibition for Hypermutant Glioblastoma Multiforme Resulting From Germline Biallelic Mismatch Repair Deficiency. J Clin Oncol. 2016 Jul 1;34(19):2206-11.
72. Schenker M, Burotto M, Richardet M, et al. Randomized, open-label, phase 2 study of nivolumab plus ipilimumab or nivolumab monotherapy in patients with advanced or metastatic solid tumors of high tumor mutational burden. J Immunother Cancer 2024; 12:e008872
73. Mei MG, Lee HJ, Palmer J, et al. Response-adapted anti-PD-1-based salvage therapy for Hodgkin lymphoma with nivolumab alone or in combination with ICE. Blood. 2022 Jun 23;139(25):3605-3616. Doi: 10.1182/blood.2022015423.
74. Zinzani P, Santoro A, Gritti G, et al. Nivolumab Combined With Brentuximab Vedotin for Relapsed/Refractory Primary Mediastinal Large B-Cell Lymphoma: Efficacy and Safety From the Phase II CheckMate 436 Study. J Clin Oncol. 2019 Nov 20;37(33):3081-3089. Doi: 10.1200/JCO.19.01492. Epub 2019 Aug 9.
75. Davis KL, Fox E, Merchant MS, et al. Nivolumab in children and young adults with relapsed or refractory solid tumours or lymphoma (ADVL1412): a multicentre, open-label, single-arm, phase 1–2 trial. The Lancet Oncology. Volume 21, issue 4, p541-550, April 01, 2020 https://doi.org/10.1016/S1470-2045(20)30023-1.
76. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Pediatric Aggressive Mature B-Cell Lymphomas. Version 1.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
77. Younes A, Santoro A, Shipp M, et al. Nivolumab for classical Hodgkin's lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: a multicentre, multicohort, single-arm phase 2 trial. Lancet Oncol. 2016 Sep;17(9):1283-94. Doi: 10.1016/S1470-2045(16)30167-X.
78. Chung C, Li J, Steuer C, et al. Phase II Multi-institutional Clinical Trial Result of Concurrent Cetuximab and Nivolumab in Recurrent and/or Metastatic Head and Neck Squamous Cell Carcinoma. Clin Cancer Res. 2022 Jun 1;28(11):2329-2338. Doi: 10.1158/1078-0432.CCR-21-3849.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

79. Zer A, Icht O, Yosef L, et al. Phase II single-arm study of nivolumab and ipilimumab (Nivo/Ipi) in previously treated classical Kaposi sarcoma (cKS). Annals of Oncology. Volume 33, Issue 7, July 2022, Pages 720-727. https://doi.org/10.1016/j.annonc.2022.03.012.
80. Pelster MS, Gruschkus SK, Bassett R, et al. Nivolumab and Ipilimumab in Metastatic Uveal Melanoma: Results From a Single-Arm Phase II Study. J Clin Oncol. 2021 Feb 20;39(6):599-607. Doi: 10.1200/JCO.20.00605.
81. Baas P, Scherpereel A, Nowak AK, et al. First-line nivolumab plus ipilimumab in unresectable malignant pleural mesothelioma (CheckMate 743): a multicentre, randomised, open-label, phase 3 trial. Lancet. 2021 Jan 30;397(10272):375-386. Doi: 10.1016/S0140-6736(20)32714-8.
82. Blank CU, Rozeman EA, Fanchi LF, et al. Neoadjuvant versus adjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma. Nat Med. 2018 Nov;24(11):1655-1661. Doi: 10.1038/s41591-018-0198-0.
83. Glutsch V, Kneitz, Gesierich A, et al. Activity of ipilimumab plus nivolumab in avelumab-refractory Merkel cell carcinoma. Cancer Immunology, Immunotherapy volume 70, pages2087–2093 (2021)
84. Wagner M, Othus M, Patel S, et al. Multicenter phase II trial (SWOG S1609, cohort 51) of ipilimumab and nivolumab in metastatic or unresectable angiosarcoma: a substudy of dual anti-CTLA-4 and anti-PD-1 blockade in rare tumors (DART). J Immunother Cancer. 2021 Aug;9(8):e002990. Doi: 10.1136/jitc-2021-002990.
85. Kim S, Wuthrick E, Blakaj D, et al. Combined nivolumab and ipilimumab with or without stereotactic body radiation therapy for advanced Merkel cell carcinoma: a randomized, open label, phase 2 trial. The Lancet. Published: September 11, 2022. Doi:https://doi.org/10.1016/S0140-6736(22)01659-2. PlumX Metrics
86. Yau T, Park JW, Finn RS, et al. Nivolumab versus sorafenib in advanced hepatocellular carcinoma (CheckMate 459): a randomised, multicentre, open-label, phase 3 trial. Lancet Oncol. 2022 Jan;23(1):77-90.
87. Kudo M, Matilla A, Santoro A, et al. CheckMate 040 cohort 5: A phase I/II study of nivolumab in patients with advanced hepatocellular carcinoma and Child-Pugh B cirrhosis. J Hepatol. 2021 Sep;75(3):600-609.
88. Long GV, Del Vecchio M, Weber J, et al. (2023). Adjuvant therapy with nivolumab versus placebo in patients with resected stage IIB/C melanoma (CheckMate 76K). SKIN The Journal of Cutaneous Medicine, 7(2), s163. https://doi.org/10.25251/skin.7.supp.163.
89. Advani RH, Moskowitz AJ, Bartlett NL, et al. Brentuximab vedotin in combination with nivolumab in relapsed or refractory Hodgkin lymphoma: 3-year study results. Blood. 2021 Aug 12;138(6):427-438. Doi: 10.1182/blood.2020009178.
90. Dagogo-Jack I, Madison RW, Lennerz JK, et al. Molecular characterization of mesothelioma: Impact of histologic type and site of origin on molecular landscape. JCO Precis Oncol 2022;6:e2100422.
91. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Colon Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.

92. van der Heijden, MS, Sonpavde G, Powles T, et al; CheckMate 901 Trial Investigators. Nivolumab plus Gemcitabine-Cisplatin in Advanced Urothelial Carcinoma. N Engl J Med. 2023 Nov 9;389(19):1778-1789. doi: 10.1056/NEJMoa2309863. Epub 2023 Oct 22. PMID: 37870949.
93. Amaria R, Reddy S, Tawbi H, et al. Neoadjuvant Immune Checkpoint Blockade in High-Risk Resectable Melanoma. Nat Med. 2018 Nov; 24(11): 1649–1654. Published online 2018 Oct 8. Doi: 10.1038/s41591-018-0197-1
94. Ma D, Ding X, Zhang C, et al. Combined targeted therapy and immunotherapy in anaplastic thyroid carcinoma with distant metastasis: A case report. World J Clin Cases 2022;10:3849-3855.
95. Kollipara R, Schneider K, Radovich M, et al. Exceptional response with immunotherapy in a patient with anaplastic thyroid cancer. Oncologist 2017;22:1149-1151.
96. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Thyroid Carcinoma. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
97. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Vaginal Cancer. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
98. Reijers ILM, Menzies AM, van Akkooi ACJ, et al. Personalized response-directed surgery and adjuvant therapy after neoadjuvant ipilimumab and nivolumab in high-risk stage III melanoma: the PRADO trial. Nat Med 2022;28:1178-1188.
99. Versluis JM, Menzies AM, Sikorska K, et al. Survival update of neoadjuvant ipilimumab plus nivolumab in macroscopic stage III melanoma in the OpACIN and OpACINneo trials. Ann Oncol 2023;34:420-430.
100. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Melanoma: Cutaneous Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
101. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab followed by Nivolumab: Colon Cancer Chemotherapy Order Template, COL68. National Comprehensive Cancer Network, 2026. NATIONAL

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.

102. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab followed by Nivolumab: Rectal Cancer Chemotherapy Order Template, REC80. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
103. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for mFOLFOX6 (Continuous Infusion Fluorouracil/Leucovorin/OXALIplatin) + Nivolumab: Gastric Cancer Chemotherapy Order Template, GAS95. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
104. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for CapeOX (Capecitabine/OXALIplatin) + Nivolumab: Gastric Cancer Chemotherapy Order Template, GAS96. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
105. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab followed by nivolumab: Ampullary Adenocarcinoma Chemotherapy Order Template, AMP22. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
106. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Mesothelioma: Peritoneal Chemotherapy Order Template, MPEM10. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
107. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab followed by Nivolumab: Central Nervous System Cancers Chemotherapy Order Template, CNS61. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

108. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Central Nervous System Cancers Chemotherapy Order Template, CNS63. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
109. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab + Ipilimumab: Non-Small Cell Lung Cancer Chemotherapy Order Template, NSC97. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
110. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: T-Cell Lymphomas Chemotherapy Order Template, TCL39. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
111. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Small Cell Lung Cancer Chemotherapy Order Template, SCL24. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
112. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Endometrial Carcinoma Chemotherapy Order Template, UTE34. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
113. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Cervical Cancer Chemotherapy Order Template, CRV35. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
114. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Vulvar Cancer Chemotherapy Order Template, VUL17. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

115. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Nivolumab: Vaginal Cancer Chemotherapy Order Template, VAG34. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
116. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Brentuximab vedotin + Nivolumab: Hodgkin Lymphoma Chemotherapy Order Template, HDL53. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
117. Herrera AF, LeBlanc M, Castellino SM, et al. Nivolumab+AVD in advanced-stage classic Hodgkin's lymphoma. N Engl J Med 2024;391:1379-1389.
118. Bröckelmann PJ, Buhen I, Meissner J et al. Nivolumab and Doxorubicin, Vinblastine, and Dacarbazine in Early-Stage Unfavorable Hodgkin Lymphoma: Final Analysis of the Randomized German Hodgkin Study Group Phase II NIVAHL Trial. J Clin Oncol. 2023 Feb 20;41(6):1193-1199. doi: 10.1200/JCO.22.02355. Epub 2022 Dec 12. PMID: 36508302.
119. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for AVD (DOXOrubicin/VinBLAStine/Dacarbazine) + Nivolumab: Hodgkin Lymphoma Chemotherapy Order Template, HDL75. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
120. Cascone T, Awad MM, Spicer JD, et al. Perioperative Nivolumab in Resectable Lung Cancer. N Engl J Med. 2024 May 16;390(19):1756-1769. doi: 10.1056/NEJMoa2311926. PMID: 38749033.
121. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Kaposi Sarcoma Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
122. Referenced with permission from the NCCN Clinical Practice Guidelines (NCCN Guidelines®) Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

123. Younes A, Brody J, Carpio C, et al. Safety and activity of ibrutinib in combination with nivolumab in patients with relapsed non-Hodgkin lymphoma or chronic lymphocytic leukaemia: a phase 1/2a study. Lancet Haematol 2019;6:e67-e78.
124. Lebbe C, Meyer N, Mortier L, et al. Initial results from a phase IIIb/IV study evaluating two dosing regimens of nivolumab (NIVO) in combination with ipilimumab (IPI) in patients with advanced melanoma (CheckMate 511)(abstract). Ann Oncol 2018;29:LBA47.
125. Andre T, Elez E, Van Cutsem E, et al; CheckMate 8HW Investigators. Nivolumab plus Ipilimumab in Microsatellite-Instability-High Metastatic Colorectal Cancer. N Engl J Med. 2024 Nov 28;391(21):2014-2026. doi: 10.1056/NEJMoa2402141. PMID: 39602630.
126. Yau T, Galle PR, Decaens T, et al. Nivolumab plus ipilimumab versus lenvatinib or sorafenib as first-line treatment for unresectable hepatocellular carcinoma (CheckMate 9DW): An open-label, randomised, phase 3 trial. Lancet 2025;405:1851-1864.
127. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Ipilimumab + Nivolumab: Hepatocellular Carcinoma Chemotherapy Order Template, HEP33. National Comprehensive Cancer Network, 2026. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Guidelines, go online to NCCN.org. Accessed July 2026.
128. Patel SP, Othus M, Chae YK, et al. Phase II basket trial of dual anti-CTLA-4 and anti-PD-1 blockade in rare tumors (DART) SWOG S1609: Adrenocortical carcinoma cohort. J Immunother Cancer 2024;12:e009074.
129. Munhoz RR, Nader-Marta G, de Camargo VP, et al. A phase 2 study of first-line nivolumab in patients with locally advanced or metastatic cutaneous squamous-cell carcinoma. *Cancer*. 2022;128(24):4223-4231. doi:10.1002/cncr.34463
130. Lang R, Welponer T, Richtig E, et al. Nivolumab for locally advanced and metastatic cutaneous squamous cell carcinoma (NIVOSQUACS study)-Phase II data covering impact of concomitant haematological malignancies. *J Eur Acad Dermatol Venereol*. 2023;37(9):1799-1810. doi:10.1111/jdv.19218
131. Harker-Murray P, Mauz-Korholz C, Leblanc T, et al. Nivolumab and brentuximab vedotin with or without bendamustine for R/R Hodgkin lymphoma in children, adolescents, and young adults. Blood 2023;141:2075-2084.
132. Greve P, Beishuizen A, Hagleitner M, et al. Nivolumab plus brentuximab vedotin +/- bendamustine combination therapy: A safe and effective treatment in pediatric recurrent and refractory classical Hodgkin lymphoma. Front Immunol 2023;14:1229558.
133. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Appendiceal Neoplasms and Cancers. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

134. Andre T, Tougeron D, Piessen G, et al.Neoadjuvant nivolumab plus ipilimumab and adjuvant nivolumab in localized deficient mismatch repair/microsatellite instability-high gastric or esophagogastric junction adenocarcinoma: The GERCOR NEONIPIGA phase II study. J Clin Oncol 2023;41:255-265.
135. Janjigian YY, Shitara K, Moehler M, et al. First-line nivolumab plus chemotherapy versus chemotherapy alone for advanced gastric, gastro-oesophageal junction, and oesophageal adenocarcinoma (CheckMate 649): a randomised, open-label, phase 3 trial. Lancet 2021;398:27-40.
136. Oaknin A, Moore K, Meyer T, et al. Nivolumab with or without ipilimumab in patients with recurrent or metastatic cervical cancer (CheckMate 358): a phase 1-2, open-label, multicohort trial. Lancet Oncol 2024;25:588-602.
137. Palmerini E, Lopez Pousa A, Grignani G, et al. Nivolumab and sunitinib in patients with advanced bone sarcomas: a multicenter, single-arm, phase 2 trial. Cancer 2025;313:e35628.
138. Long GV, Tykodi SS, Schneider JG, et al. Assessment of nivolumab exposure and clinical safety of 480 mg every 4 weeks flat-dosing schedule in patients with cancer. Ann Oncol. 2018;29(11):2208-2213.
139. Gao B, Carlino MS, Michael M, et al. Nivolumab and Ipilimumab Combination Treatment in Advanced Ovarian and Endometrial Clear Cell Cancers: A Nonrandomized Clinical Trial. JAMA Oncol. 2025 Sep 1;11(9):982-989.
140. Chae YK, Othus M, Patel, SP, et al. A phase II basket trial of dual anti-CTLA-4 and anti-PD-1 blockade in rare tumors (DART) SWOG S1609: durable responses and delayed pseudoprogression in small cell carcinoma of the ovary, hypercalcemic type cohort. Cancer Commun (Lond). 2025 Aug;45(8):976-986.
141. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Ovarian Cancer Including Fallopian Tube Cancer and Primary Peritoneal Cancer. Version 4.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.
142. Sehgal K, Pappa T, Shin KY, et al. Dual Immune Checkpoint Inhibition in Patients With Aggressive Thyroid Carcinoma: A Phase 2 Nonrandomized Clinical Trial. JAMA Oncol. 2024 Dec 1;10(12):1663-1671. doi: 10.1001/jamaoncol.2024.4019.
143. Forde PM, Spicer JD, Provencio M, et al. Overall survival with neoadjuvant nivolumab plus chemotherapy in lung cancer. N Engl J Med 2025;393:741-752.
144. Cascone T, Awad M, Spicer J, et al. Perioperative nivolumab vs placebo in patients with resectable NSCLC: Updated survival and biomarker analyses from CheckMate 77T [abstract]. J Clin Oncol 2025;43(Suppl):Abstract LBA8010.
145. Referenced with permission from the NCCN Drugs & Biologics Compendium (NCCN Compendium®) Histiocytic Neoplasms. Version 2.2026. National Comprehensive Cancer Network, 2026. The NCCN Compendium® is a derivative work of the NCCN Guidelines®.

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, and NCCN GUIDELINES® are trademarks owned by the National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the Compendium, go online to NCCN.org. Accessed July 2026.

146. Bose S, Robles J, McCall CM, et al. Favorable response to nivolumab in a young adult patient with metastatic histiocytic sarcoma. Pediatr Blood Cancer 2019;66e27491. Available at: https://pmc.ncbi.nlm.nih.gov/articles/PMC6433376/

147. Campedel L, Karroubi D, Vozy A, et al. Malignant histiocytosis with PD-L1 expression-dramatic response to nivolumab. Mayo Clin Proc 2022;97:1401-1403. Available at: https://www.mayoclinicproceedings.org/article/S0025-6196(22)00299-3/fulltext

148. Imataki O, Uemura M, Fujita M, Kadowaki N. Application of PD-L1 blockade in refractory histiocytic sarcoma: a case report. Mol Clin Oncol 2022;17:136. Available at: https://pmc.ncbi.nlm.nih.gov/articles/PMC9353867/

Appendix A – Non-Quantitative Treatment Limitations (NQTL) Factor Checklist

Non-quantitative treatment limitations (NQTLs) refer to the methods, guidelines, standards of evidence, or other conditions that can restrict how long or to what extent benefits are provided under a health plan. These may include things like utilization review or prior authorization. The utilization management NQTL applies comparably, and not more stringently, to mental health/substance use disorder (MH/SUD) Medical Benefit Prescription Drugs and medical/surgical (M/S) Medical Benefit Prescription Drugs. The table below lists the factors that were considered in designing and applying prior authorization to this drug/drug group, and a summary of the conclusions that Prime's assessment led to for each.

Factor	Conclusion
Indication	Yes: Consider for PA
Safety and efficacy	No: PA not a priority
Potential for misuse/abuse	No: PA not a priority
Cost of drug	Yes: Consider for PA

Appendix 1 – Covered Diagnosis Codes

ICD-10	ICD-10 Description
C00.3	Malignant neoplasm of upper lip, inner aspect
C00.4	Malignant neoplasm of lower lip, inner aspect
C00.5	Malignant neoplasm of lip, unspecified, inner aspect
C00.6	Malignant neoplasm of commissure of lip, unspecified
C00.8	Malignant neoplasm of overlapping sites of lip
C00.9	Malignant neoplasm of lip, unspecified
C01	Malignant neoplasm of base of tongue
C02.0	Malignant neoplasm of dorsal surface of tongue
C02.1	Malignant neoplasm of border of tongue
C02.2	Malignant neoplasm of ventral surface of tongue

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C02.3	Malignant neoplasm of anterior two-thirds of tongue, part unspecified
C02.4	Malignant neoplasm of lingual tonsil
C02.8	Malignant neoplasm of overlapping sites of tongue
C02.9	Malignant neoplasm of tongue, unspecified
C03.0	Malignant neoplasm of upper gum
C03.1	Malignant neoplasm of lower gum
C03.9	Malignant neoplasm of gum, unspecified
C04.0	Malignant neoplasm of anterior floor of mouth
C04.1	Malignant neoplasm of lateral floor of mouth
C04.8	Malignant neoplasm of overlapping sites of floor of mouth
C04.9	Malignant neoplasm of floor of mouth, unspecified
C05.0	Malignant neoplasm of hard palate
C05.1	Malignant neoplasm of soft palate
C05.2	Malignant neoplasm of uvula
C05.8	Malignant neoplasm of overlapping sites of palate
C05.9	Malignant neoplasm of palate, unspecified
C06.0	Malignant neoplasm of cheek mucosa
C06.2	Malignant neoplasm of retromolar area
C06.80	Malignant neoplasm of overlapping sites of unspecified parts of mouth
C06.89	Malignant neoplasm of overlapping sites of other parts of mouth
C06.9	Malignant neoplasm of mouth, unspecified
C09.0	Malignant neoplasm of tonsillar fossa
C09.1	Malignant neoplasm of tonsillar pillar (anterior) (posterior)
C09.8	Malignant neoplasm of overlapping sites of tonsil
C09.9	Malignant neoplasm of tonsil, unspecified
C10.0	Malignant neoplasm of vallecula
C10.1	Malignant neoplasm of anterior surface of epiglottis
C10.2	Malignant neoplasm of lateral wall of oropharynx
C10.3	Malignant neoplasm of posterior wall of oropharynx
C10.4	Malignant neoplasm of branchial cleft
C10.8	Malignant neoplasm of overlapping sites of oropharynx
C10.9	Malignant neoplasm of oropharynx, unspecified
C11.0	Malignant neoplasm of superior wall of nasopharynx
C11.1	Malignant neoplasm of posterior wall of nasopharynx
C11.2	Malignant neoplasm of lateral wall of nasopharynx

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C11.3	Malignant neoplasm of anterior wall of nasopharynx
C11.8	Malignant neoplasm of overlapping sites of nasopharynx
C11.9	Malignant neoplasm of nasopharynx, unspecified
C12	Malignant neoplasm of pyriform sinus
C13.0	Malignant neoplasm of postcricoid region
C13.1	Malignant neoplasm of aryepiglottic fold, hypopharyngeal aspect
C13.2	Malignant neoplasm of posterior wall of hypopharynx
C13.8	Malignant neoplasm of overlapping sites of hypopharynx
C13.9	Malignant neoplasm of hypopharynx, unspecified
C14.0	Malignant neoplasm of pharynx, unspecified
C14.2	Malignant neoplasm of Waldeyer's ring
C14.8	Malignant neoplasm of overlapping sites of lip, oral cavity and pharynx
C15.3	Malignant neoplasm of upper third of esophagus
C15.4	Malignant neoplasm of middle third of esophagus
C15.5	Malignant neoplasm of lower third of esophagus
C15.8	Malignant neoplasm of overlapping sites of esophagus
C15.9	Malignant neoplasm of esophagus, unspecified
C16.0	Malignant neoplasm of cardia
C16.1	Malignant neoplasm of fundus of stomach
C16.2	Malignant neoplasm of body of stomach
C16.3	Malignant neoplasm of pyloric antrum
C16.4	Malignant neoplasm of pylorus
C16.5	Malignant neoplasm of lesser curvature of stomach, unspecified
C16.6	Malignant neoplasm of greater curvature of stomach, unspecified
C16.8	Malignant neoplasm of overlapping sites of stomach
C16.9	Malignant neoplasm of stomach, unspecified
C17.0	Malignant neoplasm of duodenum
C17.1	Malignant neoplasm of jejunum
C17.2	Malignant neoplasm of ileum
C17.3	Meckel's diverticulum, malignant
C17.8	Malignant neoplasm of overlapping sites of small intestine
C17.9	Malignant neoplasm of small intestine, unspecified
C18.0	Malignant neoplasm of cecum
C18.1	Malignant neoplasm of appendix
C18.2	Malignant neoplasm of ascending colon

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C18.3	Malignant neoplasm of hepatic flexure
C18.4	Malignant neoplasm of transverse colon
C18.5	Malignant neoplasm of splenic flexure
C18.6	Malignant neoplasm of descending colon
C18.7	Malignant neoplasm of sigmoid colon
C18.8	Malignant neoplasm of overlapping sites of colon
C18.9	Malignant neoplasm of colon, unspecified
C19	Malignant neoplasm of rectosigmoid junction
C20	Malignant neoplasm of rectum
C21.0	Malignant neoplasm of anus, unspecified
C21.1	Malignant neoplasm of anal canal
C21.2	Malignant neoplasm of cloacogenic zone
C21.8	Malignant neoplasm of overlapping sites of rectum, anus and anal canal
C22.0	Liver cell carcinoma
C22.1	Intrahepatic bile duct carcinoma
C22.3	Angiosarcoma of liver
C22.8	Malignant neoplasm of liver, primary, unspecified as to type
C22.9	Malignant neoplasm of liver, not specified as primary or secondary
C23	Malignant neoplasm of gallbladder
C24.0	Malignant neoplasm of extrahepatic bile duct
C24.1	Malignant neoplasm of ampulla of Vater
C24.8	Malignant neoplasm of overlapping sites of biliary tract
C24.9	Malignant neoplasm of biliary tract, unspecified
C30.0	Malignant neoplasm of nasal cavity
C31.0	Malignant neoplasm of maxillary sinus
C31.1	Malignant neoplasm of ethmoidal sinus
C32.0	Malignant neoplasm of glottis
C32.1	Malignant neoplasm of supraglottis
C32.2	Malignant neoplasm of subglottis
C32.3	Malignant neoplasm of laryngeal cartilage
C32.8	Malignant neoplasm of overlapping sites of larynx
C32.9	Malignant neoplasm of larynx, unspecified
C33	Malignant neoplasm of trachea
C34.00	Malignant neoplasm of unspecified main bronchus
C34.01	Malignant neoplasm of right main bronchus

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C34.02	Malignant neoplasm of left main bronchus
C34.10	Malignant neoplasm of upper lobe, unspecified bronchus or lung
C34.11	Malignant neoplasm of upper lobe, right bronchus or lung
C34.12	Malignant neoplasm of upper lobe, left bronchus or lung
C34.2	Malignant neoplasm of middle lobe, bronchus or lung
C34.30	Malignant neoplasm of lower lobe, unspecified bronchus or lung
C34.31	Malignant neoplasm of lower lobe, right bronchus or lung
C34.32	Malignant neoplasm of lower lobe, left bronchus or lung
C34.80	Malignant neoplasm of overlapping sites of unspecified bronchus and lung
C34.81	Malignant neoplasm of overlapping sites of right bronchus and lung
C34.82	Malignant neoplasm of overlapping sites of left bronchus and lung
C34.90	Malignant neoplasm of unspecified part of unspecified bronchus or lung
C34.91	Malignant neoplasm of unspecified part of right bronchus or lung
C34.92	Malignant neoplasm of unspecified part of left bronchus or lung
C40.00	Malignant neoplasm of scapula and long bones of unspecified upper limb
C40.01	Malignant neoplasm of scapula and long bones of right upper limb
C40.02	Malignant neoplasm of scapula and long bones of left upper limb
C40.10	Malignant neoplasm of short bones of unspecified upper limb
C40.11	Malignant neoplasm of short bones of right upper limb
C40.12	Malignant neoplasm of short bones of left upper limb
C40.20	Malignant neoplasm of long bones of unspecified lower limb
C40.21	Malignant neoplasm of long bones of right lower limb
C40.22	Malignant neoplasm of long bones of left lower limb
C40.30	Malignant neoplasm of short bones of unspecified lower limb
C40.31	Malignant neoplasm of short bones of right lower limb
C40.32	Malignant neoplasm of short bones of left lower limb
C40.80	Malignant neoplasm of overlapping sites of bone and articular cartilage of unspecified limb
C40.81	Malignant neoplasm of overlapping sites of bone and articular cartilage of right limb
C40.82	Malignant neoplasm of overlapping sites of bone and articular cartilage of left limb
C40.90	Malignant neoplasm of unspecified bones and articular cartilage of unspecified limb
C40.91	Malignant neoplasm of unspecified bones and articular cartilage of right limb
C40.92	Malignant neoplasm of unspecified bones and articular cartilage of left limb
C41.0	Malignant neoplasm of bones of skull and face
C41.1	Malignant neoplasm of mandible
C41.2	Malignant neoplasm of vertebral column

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C41.3	Malignant neoplasm of ribs, sternum and clavicle
C41.4	Malignant neoplasm of pelvic bones, sacrum and coccyx
C41.9	Malignant neoplasm of bone and articular cartilage, unspecified
C43.0	Malignant melanoma of lip
C43.10	Malignant melanoma of unspecified eyelid, including canthus
C43.111	Malignant melanoma of right upper eyelid, including canthus
C43.112	Malignant melanoma of right lower eyelid, including canthus
C43.121	Malignant melanoma of left upper eyelid, including canthus
C43.122	Malignant melanoma of left lower eyelid, including canthus
C43.20	Malignant melanoma of unspecified ear and external auricular canal
C43.21	Malignant melanoma of right ear and external auricular canal
C43.22	Malignant melanoma of left ear and external auricular canal
C43.30	Malignant melanoma of unspecified part of face
C43.31	Malignant melanoma of nose
C43.39	Malignant melanoma of other parts of face
C43.4	Malignant melanoma of scalp and neck
C43.51	Malignant melanoma of anal skin
C43.52	Malignant melanoma of skin of breast
C43.59	Malignant melanoma of other part of trunk
C43.60	Malignant melanoma of unspecified upper limb, including shoulder
C43.61	Malignant melanoma of right upper limb, including shoulder
C43.62	Malignant melanoma of left upper limb, including shoulder
C43.70	Malignant melanoma of unspecified lower limb, including hip
C43.71	Malignant melanoma of right lower limb, including hip
C43.72	Malignant melanoma of left lower limb, including hip
C43.8	Malignant melanoma of overlapping sites of skin
C43.9	Malignant melanoma of skin, unspecified
C44.02	Squamous cell carcinoma of skin of lip
C44.121	Squamous cell carcinoma of skin of unspecified eyelid, including canthus
C44.1221	Squamous cell carcinoma of skin of right upper eyelid, including canthus
C44.1222	Squamous cell carcinoma of skin of right lower eyelid, including canthus
C44.1291	Squamous cell of carcinoma of skin of left upper eyelid, including canthus
C44.1292	Squamous cell carcinoma of skin of left lower eyelid, including canthus
C44.221	Squamous cell carcinoma of skin of unspecified ear and external auricular canal
C44.222	Squamous cell carcinoma of skin of right ear and external auricular canal

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C44.229	Squamous cell carcinoma of skin of left ear and external auricular canal
C44.320	Squamous cell carcinoma of skin of unspecified parts of face
C44.321	Squamous cell carcinoma of skin of nose
C44.329	Squamous cell carcinoma of skin of other parts of face
C44.42	Squamous cell carcinoma of skin of scalp and neck
C44.520	Squamous cell carcinoma of anal skin
C44.521	Squamous cell carcinoma of skin of breast
C44.529	Squamous cell carcinoma of skin of other part of trunk
C44.621	Squamous cell carcinoma of skin of unspecified upper limb, including shoulder
C44.622	Squamous cell carcinoma of skin of right upper limb, including shoulder
C44.629	Squamous cell carcinoma of skin of left upper limb, including shoulder
C44.721	Squamous cell carcinoma of skin of unspecified lower limb, including hip
C44.722	Squamous cell carcinoma of skin of right lower limb, including hip
C44.729	Squamous cell carcinoma of skin of left lower limb, including hip
C44.82	Squamous cell carcinoma of overlapping sites of skin
C44.92	Squamous cell carcinoma of skin, unspecified
C45.0	Mesothelioma of pleura
C45.1	Mesothelioma of peritoneum
C45.2	Mesothelioma of pericardium
C45.7	Mesothelioma of other sites
C45.9	Mesothelioma, unspecified
C46.0	Kaposi’s sarcoma of skin
C46.1	Kaposi’s sarcoma of soft tissue
C46.2	Kaposi’s sarcoma of palate
C46.3	Kaposi’s sarcoma of lymph nodes
C46.4	Kaposi’s sarcoma of gastrointestinal sites
C46.50	Kaposi’s sarcoma of unspecified lung
C46.51	Kaposi’s sarcoma of right lung
C46.52	Kaposi’s sarcoma of left lung
C46.7	Kaposi’s sarcoma of other sites
C46.9	Kaposi’s sarcoma, unspecified
C47.0	Malignant neoplasm of peripheral nerves of head, face and neck
C47.10	Malignant neoplasm of peripheral nerves of unspecified upper limb, including shoulder
C47.11	Malignant neoplasm of peripheral nerves of right upper limb, including shoulder
C47.12	Malignant neoplasm of peripheral nerves of left upper limb, including shoulder
C47.20	Malignant neoplasm of peripheral nerves of unspecified lower limb, including hip

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C47.21	Malignant neoplasm of peripheral nerves of right lower limb, including hip
C47.22	Malignant neoplasm of peripheral nerves of left lower limb, including hip
C47.3	Malignant neoplasm of peripheral nerves of thorax
C47.4	Malignant neoplasm of peripheral nerves of abdomen
C47.5	Malignant neoplasm of peripheral nerves of pelvis
C47.6	Malignant neoplasm of peripheral nerves of trunk, unspecified
C47.8	Malignant neoplasm of overlapping sites of peripheral nerves and autonomic nervous system
C47.9	Malignant neoplasm of peripheral nerves and autonomic nervous system, unspecified
C48.0	Malignant neoplasm of retroperitoneum
C48.1	Malignant neoplasm of specified parts of peritoneum
C48.2	Malignant neoplasm of peritoneum, unspecified
C48.8	Malignant neoplasm of overlapping sites of retroperitoneum and peritoneum
C49.0	Malignant neoplasm of connective and soft tissue of head, face and neck
C49.10	Malignant neoplasm of connective and soft tissue of unspecified upper limb, including shoulder
C49.11	Malignant neoplasm of connective and soft tissue of right upper limb, including shoulder
C49.12	Malignant neoplasm of connective and soft tissue of left upper limb, including shoulder
C49.20	Malignant neoplasm of connective and soft tissue of unspecified lower limb, including hip
C49.21	Malignant neoplasm of connective and soft tissue of right lower limb, including hip
C49.22	Malignant neoplasm of connective and soft tissue of left lower limb, including hip
C49.3	Malignant neoplasm of connective and soft tissue of thorax
C49.4	Malignant neoplasm of connective and soft tissue of abdomen
C49.5	Malignant neoplasm of connective and soft tissue of pelvis
C49.6	Malignant neoplasm of connective and soft tissue of trunk, unspecified
C49.8	Malignant neoplasm of overlapping sites of connective and soft tissue
C49.9	Malignant neoplasm of connective and soft tissue, unspecified
C4A.0	Merkel cell carcinoma of lip
C4A.10	Merkel cell carcinoma of eyelid, including canthus
C4A.111	Merkel cell carcinoma of right upper eyelid, including canthus
C4A.112	Merkel cell carcinoma of right lower eyelid, including canthus
C4A.121	Merkel cell carcinoma of left upper eyelid, including canthus
C4A.122	Merkel cell carcinoma of left lower eyelid, including canthus
C4A.20	Merkel cell carcinoma of unspecified ear and external auricular canal
C4A.21	Merkel cell carcinoma of right ear and external auricular canal
C4A.22	Merkel cell carcinoma of left ear and external auricular canal
C4A.30	Merkel cell carcinoma of unspecified part of face
C4A.31	Merkel cell carcinoma of nose
C4A.39	Merkel cell carcinoma of other parts of face

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C4A.4	Merkel cell carcinoma of scalp and neck
C4A.51	Merkel cell carcinoma of anal skin
C4A.52	Merkel cell carcinoma of skin of breast
C4A.59	Merkel cell carcinoma of other part of trunk
C4A.60	Merkel cell carcinoma of unspecified upper limb, including shoulder
C4A.61	Merkel cell carcinoma of right upper limb, including shoulder
C4A.62	Merkel cell carcinoma of left upper limb, including shoulder
C4A.70	Merkel cell carcinoma of unspecified lower limb, including hip
C4A.71	Merkel cell carcinoma of right lower limb, including hip
C4A.72	Merkel cell carcinoma of left lower limb, including hip
C4A.8	Merkel cell carcinoma of overlapping sites
C4A.9	Merkel cell carcinoma, unspecified
C51.0	Malignant neoplasm of labium majus
C51.1	Malignant neoplasm of labium minus
C51.2	Malignant neoplasm of clitoris
C51.8	Malignant neoplasm of overlapping sites of vulva
C51.9	Malignant neoplasm of vulva, unspecified
C52	Malignant neoplasm of vagina
C53.0	Malignant neoplasm of endocervix
C53.1	Malignant neoplasm of exocervix
C53.8	Malignant neoplasm of overlapping sites of cervix uteri
C53.9	Malignant neoplasm of cervix uteri, unspecified
C54.0	Malignant neoplasm of isthmus uteri
C54.1	Malignant neoplasm of endometrium
C54.2	Malignant neoplasm of myometrium
C54.3	Malignant neoplasm of fundus uteri
C54.8	Malignant neoplasm of overlapping sites of corpus uteri
C54.9	Malignant neoplasm of corpus uteri, unspecified
C55	Malignant neoplasm of uterus, part unspecified
C56.1	Malignant neoplasm of right ovary
C56.2	Malignant neoplasm of left ovary
C56.3	Malignant neoplasm of bilateral ovaries
C56.9	Malignant neoplasm of unspecified ovary
C57.00	Malignant neoplasm of unspecified fallopian tube
C57.01	Malignant neoplasm of right fallopian tube
C57.02	Malignant neoplasm of left fallopian tube
C57.10	Malignant neoplasm of unspecified broad ligament

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C57.11	Malignant neoplasm of right broad ligament
C57.12	Malignant neoplasm of left broad ligament
C57.20	Malignant neoplasm of unspecified round ligament
C57.21	Malignant neoplasm of right round ligament
C57.22	Malignant neoplasm of left round ligament
C57.3	Malignant neoplasm of parametrium
C57.4	Malignant neoplasm of uterine adnexa, unspecified
C57.7	Malignant neoplasm of other specified female genital organs
C57.8	Malignant neoplasm of overlapping sites of female genital organs
C57.9	Malignant neoplasm of female genital organ, unspecified
C58	Malignant neoplasm of placenta
C61	Malignant neoplasm of prostate
C64.1	Malignant neoplasm of right kidney, except renal pelvis
C64.2	Malignant neoplasm of left kidney, except renal pelvis
C64.9	Malignant neoplasm of unspecified kidney, except renal pelvis
C65.1	Malignant neoplasm of right renal pelvis
C65.2	Malignant neoplasm of left renal pelvis
C65.9	Malignant neoplasm of unspecified renal pelvis
C66.1	Malignant neoplasm of right ureter
C66.2	Malignant neoplasm of left ureter
C66.9	Malignant neoplasm of unspecified ureter
C67.0	Malignant neoplasm of trigone of bladder
C67.1	Malignant neoplasm of dome of bladder
C67.2	Malignant neoplasm of lateral wall of bladder
C67.3	Malignant neoplasm of anterior wall of bladder
C67.4	Malignant neoplasm of posterior wall of bladder
C67.5	Malignant neoplasm of bladder neck
C67.6	Malignant neoplasm of ureteric orifice
C67.7	Malignant neoplasm of urachus
C67.8	Malignant neoplasm of overlapping sites of bladder
C67.9	Malignant neoplasm of bladder, unspecified
C68.0	Malignant neoplasm of urethra
C69.30	Malignant neoplasm of unspecified choroid
C69.31	Malignant neoplasm of right choroid
C69.32	Malignant neoplasm of left choroid
C69.40	Malignant neoplasm of unspecified ciliary body

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C69.41	Malignant neoplasm of right ciliary body
C69.42	Malignant neoplasm of left ciliary body
C69.60	Malignant neoplasm of unspecified orbit
C69.61	Malignant neoplasm of right orbit
C69.62	Malignant neoplasm of left orbit
C71.0	Malignant neoplasm of cerebrum, except lobes and ventricles
C71.1	Malignant neoplasm of frontal lobe
C71.2	Malignant neoplasm of temporal lobe
C71.3	Malignant neoplasm of parietal lobe
C71.4	Malignant neoplasm of occipital lobe
C71.5	Malignant neoplasm of cerebral ventricle
C71.6	Malignant neoplasm of cerebellum
C71.7	Malignant neoplasm of brain stem
C71.8	Malignant neoplasm of overlapping sites of brain
C71.9	Malignant neoplasm of brain, unspecified
C72.0	Malignant neoplasm of spinal cord
C72.1	Malignant neoplasm of cauda equina
C72.9	Malignant neoplasm of central nervous system, unspecified
C73	Malignant neoplasm of thyroid gland
C74.00	Malignant neoplasm of cortex of unspecified adrenal gland
C74.01	Malignant neoplasm of cortex of right adrenal gland
C74.02	Malignant neoplasm of cortex of left adrenal gland
C74.90	Malignant neoplasm of unspecified part of unspecified adrenal gland
C74.91	Malignant neoplasm of unspecified part of right adrenal gland
C74.92	Malignant neoplasm of unspecified part of left adrenal gland
C76.0	Malignant neoplasm of head, face and neck
C78.00	Secondary malignant neoplasm of unspecified lung
C78.01	Secondary malignant neoplasm of right lung
C78.02	Secondary malignant neoplasm of left lung
C78.6	Secondary malignant neoplasm of retroperitoneum and peritoneum
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C79.31	Secondary malignant neoplasm of brain
C79.70	Secondary malignant neoplasm of unspecified adrenal gland
C79.71	Secondary malignant neoplasm of right adrenal gland
C79.72	Secondary malignant neoplasm of left adrenal gland

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C7A.1	Malignant poorly differentiated neuroendocrine tumors
C7A.8	Other malignant neuroendocrine tumors
C7B.00	Secondary carcinoid tumors, unspecified site
C7B.01	Secondary carcinoid tumors of distant lymph nodes
C7B.02	Secondary carcinoid tumors of liver
C7B.03	Secondary carcinoid tumors of bone
C7B.04	Secondary carcinoid tumors of peritoneum
C7B.1	Secondary Merkel cell carcinoma
C7B.8	Other secondary neuroendocrine tumors
C81.10	Nodular sclerosis Hodgkin lymphoma, unspecified site
C81.11	Nodular sclerosis Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.12	Nodular sclerosis Hodgkin lymphoma, intrathoracic lymph nodes
C81.13	Nodular sclerosis Hodgkin lymphoma, intra-abdominal lymph nodes
C81.14	Nodular sclerosis Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.15	Nodular sclerosis Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.16	Nodular sclerosis Hodgkin lymphoma, intrapelvic lymph nodes
C81.17	Nodular sclerosis Hodgkin lymphoma, spleen
C81.18	Nodular sclerosis Hodgkin lymphoma, lymph nodes of multiple sites
C81.19	Nodular sclerosis Hodgkin lymphoma, extranodal and solid organ sites
C81.20	Mixed cellularity Hodgkin lymphoma, unspecified site
C81.21	Mixed cellularity Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.22	Mixed cellularity Hodgkin lymphoma, intrathoracic lymph nodes
C81.23	Mixed cellularity Hodgkin lymphoma, intra-abdominal lymph nodes
C81.24	Mixed cellularity Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.25	Mixed cellularity Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.26	Mixed cellularity Hodgkin lymphoma, intrapelvic lymph nodes
C81.27	Mixed cellularity Hodgkin lymphoma, spleen
C81.28	Mixed cellularity Hodgkin lymphoma, lymph nodes of multiple sites
C81.29	Mixed cellularity Hodgkin lymphoma, extranodal and solid organ sites
C81.30	Lymphocyte depleted Hodgkin lymphoma, unspecified site
C81.31	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.32	Lymphocyte depleted Hodgkin lymphoma, intrathoracic lymph nodes
C81.33	Lymphocyte depleted Hodgkin lymphoma, intra-abdominal lymph nodes
C81.34	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.35	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of inguinal region and lower limb

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C81.36	Lymphocyte depleted Hodgkin lymphoma, intrapelvic lymph nodes
C81.37	Lymphocyte depleted Hodgkin lymphoma, spleen
C81.38	Lymphocyte depleted Hodgkin lymphoma, lymph nodes of multiple sites
C81.39	Lymphocyte depleted Hodgkin lymphoma, extranodal and solid organ sites
C81.40	Lymphocyte-rich Hodgkin lymphoma, unspecified site
C81.41	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of head, face, and neck
C81.42	Lymphocyte-rich Hodgkin lymphoma, intrathoracic lymph nodes
C81.43	Lymphocyte-rich Hodgkin lymphoma, intra-abdominal lymph nodes
C81.44	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of axilla and upper limb
C81.45	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of inguinal region and lower limb
C81.46	Lymphocyte-rich Hodgkin lymphoma, intrapelvic lymph nodes
C81.47	Lymphocyte-rich Hodgkin lymphoma, spleen
C81.48	Lymphocyte-rich Hodgkin lymphoma, lymph nodes of multiple sites
C81.49	Lymphocyte-rich Hodgkin lymphoma, extranodal and solid organ sites
C81.70	Other Hodgkin lymphoma unspecified site
C81.71	Other Hodgkin lymphoma lymph nodes of head, face, and neck
C81.72	Other Hodgkin lymphoma intrathoracic lymph nodes
C81.73	Other Hodgkin lymphoma intra-abdominal lymph nodes
C81.74	Other Hodgkin lymphoma lymph nodes of axilla and upper limb
C81.75	Other Hodgkin lymphoma lymph nodes of inguinal region and lower limb
C81.76	Other Hodgkin lymphoma intrapelvic lymph nodes
C81.77	Other Hodgkin lymphoma spleen
C81.78	Other Hodgkin lymphoma lymph nodes of multiple sites
C81.79	Other Hodgkin lymphoma extranodal and solid organ sites
C81.90	Hodgkin lymphoma, unspecified site
C81.91	Hodgkin lymphoma, unspecified lymph nodes of head, face, and neck
C81.92	Hodgkin lymphoma, unspecified intrathoracic lymph nodes
C81.93	Hodgkin lymphoma, unspecified intra-abdominal lymph nodes
C81.94	Hodgkin lymphoma, unspecified lymph nodes of axilla and upper limb
C81.95	Hodgkin lymphoma, unspecified lymph nodes of inguinal region and lower limb
C81.96	Hodgkin lymphoma, unspecified intrapelvic lymph nodes
C81.97	Hodgkin lymphoma, unspecified spleen
C81.98	Hodgkin lymphoma, unspecified lymph nodes of multiple sites
C81.99	Hodgkin lymphoma, unspecified extranodal and solid organ sites
C83.00	Small cell B-cell lymphoma, unspecified site

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C83.01	Small cell B-cell lymphoma, lymph nodes of head, face, and neck
C83.02	Small cell B-cell lymphoma, intrathoracic lymph nodes
C83.03	Small cell B-cell lymphoma, intra-abdominal lymph nodes
C83.04	Small cell B-cell lymphoma, lymph nodes of axilla and upper limb
C83.05	Small cell B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.06	Small cell B-cell lymphoma, intrapelvic lymph nodes
C83.07	Small cell B-cell lymphoma, spleen
C83.08	Small cell B-cell lymphoma, lymph nodes of multiple sites
C83.09	Small cell B-cell lymphoma, extranodal and solid organ sites
C83.30	Diffuse large B-cell lymphoma, unspecified site
C83.31	Diffuse large B-cell lymphoma, lymph nodes of head, face, and neck
C83.32	Diffuse large B-cell lymphoma, intrathoracic lymph nodes
C83.33	Diffuse large B-cell lymphoma, intra-abdominal lymph nodes
C83.34	Diffuse large B-cell lymphoma, lymph nodes of axilla and upper limb
C83.35	Diffuse large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C83.36	Diffuse large B-cell lymphoma, intrapelvic lymph nodes
C83.37	Diffuse large B-cell lymphoma, spleen
C83.38	Diffuse large B-cell lymphoma, lymph nodes of multiple sites
C83.39	Diffuse large B-cell lymphoma, extranodal and solid organ sites
C84.90	Mature T/NK-cell lymphomas, unspecified, unspecified site
C84.91	Mature T/NK-cell lymphomas, unspecified, lymph nodes of head, face, and neck
C84.92	Mature T/NK-cell lymphomas, unspecified, intrathoracic lymph nodes
C84.93	Mature T/NK-cell lymphomas, unspecified, intra-abdominal lymph nodes
C84.94	Mature T/NK-cell lymphomas, unspecified, lymph nodes of axilla and upper limb
C84.95	Mature T/NK-cell lymphomas, unspecified, lymph nodes of inguinal region and lower limb
C84.96	Mature T/NK-cell lymphomas, unspecified, intrapelvic lymph nodes
C84.97	Mature T/NK-cell lymphomas, unspecified, spleen
C84.98	Mature T/NK-cell lymphomas, unspecified, lymph nodes of multiple sites
C84.99	Mature T/NK-cell lymphomas, unspecified, extranodal and solid organ sites
C84.Z0	Other mature T/NK-cell lymphomas, unspecified site
C84.Z1	Other mature T/NK-cell lymphomas, lymph nodes of head, face, and neck
C84.Z2	Other mature T/NK-cell lymphomas, intrathoracic lymph nodes
C84.Z3	Other mature T/NK-cell lymphomas, intra-abdominal lymph nodes
C84.Z4	Other mature T/NK-cell lymphomas, lymph nodes of axilla and upper limb
C84.Z5	Other mature T/NK-cell lymphomas, lymph nodes of inguinal region and lower limb

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

C84.Z6	Other mature T/NK-cell lymphomas, intrapelvic lymph nodes
C84.Z7	Other mature T/NK-cell lymphomas, spleen
C84.Z8	Other mature T/NK-cell lymphomas, lymph nodes of multiple sites
C84.Z9	Other mature T/NK-cell lymphomas, extranodal and solid organ sites
C85.20	Mediastinal (thymic) large B-cell lymphoma, unspecified site
C85.21	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of head, face and neck
C85.22	Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes
C85.23	Mediastinal (thymic) large B-cell lymphoma, intra-abdominal lymph nodes
C85.24	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of axilla and upper limb
C85.25	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of inguinal region and lower limb
C85.26	Mediastinal (thymic) large B-cell lymphoma, intrapelvic lymph nodes
C85.27	Mediastinal (thymic) large B-cell lymphoma, spleen
C85.28	Mediastinal (thymic) large B-cell lymphoma, lymph nodes of multiple sites
C85.29	Mediastinal (thymic) large B-cell lymphoma, extranodal and solid organ sites
C86.00	Extranodal NK/T-cell lymphoma, nasal type not having achieved remission
C91.10	Chronic lymphocytic leukemia of B-cell type not having achieved remission
C91.12	Chronic lymphocytic leukemia of B-cell type in relapse
C96.20	Malignant mast cell neoplasm, unspecified
C96.29	Other malignant mast cell neoplasm
C96.9	Malignant neoplasm of lymphoid, hematopoietic and related tissue, unspecified
C96.Z	Other specified malignant neoplasms of lymphoid, hematopoietic and related tissue
D09.0	Carcinoma in situ of bladder
D37.02	Neoplasm of uncertain behavior of tongue
D37.04	Neoplasm of uncertain behavior of the minor salivary glands
D37.05	Neoplasm of uncertain behavior of pharynx
D37.09	Neoplasm of uncertain behavior of other specified sites of the oral cavity
D37.1	Neoplasm of uncertain behavior of stomach
D37.3	Neoplasm of uncertain behavior of appendix
D37.8	Neoplasm of uncertain behavior of other specified digestive organs
D37.9	Neoplasm of uncertain behavior of digestive organ, unspecified
D38.0	Neoplasm of uncertain behavior of larynx
D38.5	Neoplasm of uncertain behavior of other respiratory organs
D38.6	Neoplasm of uncertain behavior of respiratory organ, unspecified
D39.2	Neoplasm of uncertain behavior of placenta
D48.60	Neoplasm of uncertain behavior of unspecified breast

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

D48.61	Neoplasm of uncertain behavior of right breast
D48.62	Neoplasm of uncertain behavior of left breast
O01.9	Hydatidiform mole, unspecified
Z85.00	Personal history of malignant neoplasm of unspecified digestive organ
Z85.01	Personal history of malignant neoplasm of esophagus
Z85.028	Personal history of other malignant neoplasm of stomach
Z85.038	Personal history of other malignant neoplasm of large intestine
Z85.068	Personal history of other malignant neoplasm of small intestine
Z85.09	Personal history of malignant neoplasm of other digestive organs
Z85.118	Personal history of other malignant neoplasm of bronchus and lung
Z85.12	Personal history of malignant neoplasm of trachea
Z85.22	Personal history of malignant neoplasm of nasal cavities, middle ear, and accessory sinuses
Z85.42	Personal history of malignant neoplasm of other parts of uterus
Z85.43	Personal history of malignant neoplasm of ovary
Z85.51	Personal history of malignant neoplasm of bladder
Z85.59	Personal history of malignant neoplasm of other urinary tract organ
Z85.71	Personal history of Hodgkin lymphoma
Z85.810	Personal history of malignant neoplasm of tongue
Z85.818	Personal history of malignant neoplasm of other sites of lip, oral cavity, and pharynx
Z85.820	Personal history of malignant melanoma of skin
Z85.821	Personal history of Merkel cell carcinoma
Z85.830	Personal history of malignant neoplasm of bone
Z85.831	Personal history of malignant neoplasm of soft tissue
Z85.841	Personal history of malignant neoplasm of brain
Z85.848	Personal history of malignant neoplasm of other parts of nervous tissue
Z85.850	Personal history of malignant neoplasm of thyroid
Z85.858	Personal history of malignant neoplasm of other endocrine glands

Appendix 2 – Centers for Medicare and Medicaid Services (CMS)

The preceding information is intended for non-Medicare coverage determinations. Medicare coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determinations (NCDs) and/or Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. Local Coverage Articles (LCAs) may also exist for claims payment purposes or to clarify benefit eligibility under Part B for drugs which may be self-administered. The following link may be used to search for NCD, LCD, or LCA documents:

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC

https://www.cms.gov/medicare-coverage-database/search.aspx. Additional indications, including any preceding information, may be applied at the discretion of the health plan.

Medicare Part B Covered Diagnosis Codes (applicable to existing NCD/LCD/LCA): N/A

Medicare Part B Administrative Contractor (MAC) Jurisdictions		
Jurisdiction	**Applicable State/US Territory**	**Contractor**
E (1)	CA, HI, NV, AS, GU, CNMI	Noridian Healthcare Solutions, LLC
F (2 & 3)	AK, WA, OR, ID, ND, SD, MT, WY, UT, AZ	Noridian Healthcare Solutions, LLC
5	KS, NE, IA, MO	Wisconsin Physicians Service Insurance Corp (WPS)
6	MN, WI, IL	National Government Services, Inc. (NGS)
H (4 & 7)	LA, AR, MS, TX, OK, CO, NM	Novitas Solutions, Inc.
8	MI, IN	Wisconsin Physicians Service Insurance Corp (WPS)
N (9)	FL, PR, VI	First Coast Service Options, Inc.
J (10)	TN, GA, AL	Palmetto GBA
M (11)	NC, SC, WV, VA (excluding below)	Palmetto GBA
L (12)	DE, MD, PA, NJ, DC (includes Arlington & Fairfax counties and the city of Alexandria in VA)	Novitas Solutions, Inc.
K (13 & 14)	NY, CT, MA, RI, VT, ME, NH	National Government Services, Inc. (NGS)
15	KY, OH	CGS Administrators, LLC

Proprietary Information. Restricted Access – Do not disseminate or copy without approval.

©2026 Prime Therapeutics Management, LLC